

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

GROWTH HORMONE TREATMENT IN CHILDREN and ADULTS

LENGTH OF AUTHORIZATION: UP TO ONE YEAR

REVIEW CRITERIA FOR CHILDREN:

Required for Approval:

- Must have approved diagnosis with supporting documentation (if the preferred product listed below is FDA indicated, trial of the preferred product is required)

Product Name	FDA Approved Indication	Comments
PREFERRED		
Short-Acting Agents		
Genotropin [®] , Norditropin [®]	Idiopathic Short Stature, Pediatric Growth Hormone deficiency, Prader-Willi Syndrome, Short stature due to Noonan Syndrome (Norditropin only), Small for Gestational Age, Turner Syndrome	N/A
Long-Acting Agents		
Ngenla [™]	Pediatric Growth Hormone deficiency	N/A
NON-PREFERRED		
Short-Acting Agents		
Humatrope [®] , Zomacton [®]	Pediatric Growth Hormone Deficiency, Idiopathic Short Stature, Short Stature Homeobox-Containing Gene (SHOX), Small for Gestational Age, Turner Syndrome	Refer to preferred short-acting agents
Nutropin AQ [®]	Pediatric Growth failure due to chronic renal insufficiency (CRI), Growth Hormone Deficiency, Idiopathic Short Stature, Turner Syndrome	
Omnitrope [®]	Pediatric Growth Hormone Deficiency, Idiopathic Short Stature, Prader-Willi Syndrome, Small for Gestational Age, Turner Syndrome	
Saizen [®]	Pediatric Growth Hormone Deficiency	
Long-Acting Agents		
Skytrofa [®] , Sogroya [®]	Growth Hormone Deficiency	Must have documented trial and failure on 2 short-acting agents and Ngenla.

- Must be ≤ 16 years of age
- Must be prescribed by an endocrinologist, pediatric endocrinologist or pediatric nephrologist

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

Idiopathic Short Stature:
❖ **Genotropin[®], Norditropin[®]**

Height: ≥ 2.25 standard deviations (SD) below the mean for age and gender
Bone age: Minimum of one year behind chronological age
Epiphyses: Confirmation of open growth plates if age 10 and older
Diagnostic Evaluation:

- A mixed or normal response $>10\text{ng/ml}$ to two Growth Hormone provocation tests (e.g., arginine, clonidine, glucagon, insulin, or levodopa)
- Growth velocity must be less than 5 cm/year
- Other pituitary hormone deficiencies (e.g., hypothyroidism, chronic ischemic disease) have been ruled out

Pediatric Growth Hormone Deficiency (GHD):
❖ **Genotropin[®], Norditropin[®], Ngenla[™]**

Growth velocity: < 5 cm/year
Present height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Bone age: Minimum of one year behind chronological age
Epiphyses: Confirmation of open growth plates if age 10 and older
Diagnostic Evaluation (one of the following):

- **Two** subnormal responses to GH provocation tests (e.g., arginine, clonidine, glucagon, insulin and levodopa): Confirmation of stimulation test(s) with peak serum GH concentration less than 10 ng/ml; **or**
- **One** abnormal GH test is sufficient and the patient has defined CNS pathology, multiple pituitary hormone deficiency (MPHD), history of irradiation, or a genetic defect affecting the GH axis; **or**
- **One** subnormal response to a GH provocation test with peak serum GH concentration less than 10ng/ml) **plus subnormal** serum levels of insulin-like growth factor 1 (IGF-I) **and/or** insulin-like growth factor binding protein 3 (IGFBP3)

Exclusionary Conditions:

- **Idiopathic Short Stature (ISS) has been ruled out (normal birth weight and GH sufficient)**
- Other pituitary hormone deficiencies (e.g., hypothyroidism, chronic ischemic disease) have been ruled out

Prader-Willi Syndrome:
❖ **Genotropin[®], Norditropin[®]**

Height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Diagnosis: Confirmed diagnosis of Prader-Willi Syndrome (*micro-deletion in the long arm of chromosome 15 or 2 maternal chromosome 15 and no paternal chromosome 15, or nonfunctional paternal chromosome 15*)
Epiphyses: Confirmation of open growth plates if age 10 and older

Small for Gestational Age (SGA):
❖ **Genotropin[®], Norditropin[®]**

Age: Greater than 2 years old
Birth weight/length: ≥ 2 standard deviations (SD) below the mean for gestational age

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

Growth velocity: Failure to manifest catch-up growth by two years of age, defined as 2 standard deviations (SD) below the mean for age and gender **AND projected height more than 1.5 SD below the midparental height (based on growth trajectory to adulthood)**
Epiphyses: Confirmation of open growth plates if age 10 and older

Turner Syndrome:
 ❖ **Genotropin®**, **Norditropin®**

Age/Gender: Females greater than 2 years old
Height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Growth Velocity: < 5 cm/year
Bone age: Less than 14 years
Diagnosis: Confirmed diagnosis of Turner Syndrome (*peripheral blood karyotype showing a 45, XO genotype*)
Epiphyses: Confirmation of open growth plates if age 10 and older

For short stature in children with SHOX (short stature homeobox-containing gene) deficiency:
 ❖ **Humatrope®**, **Zomacton®**

Height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Growth Velocity: < 5 cm/year
Bone age: Less than 14 years
Diagnosis: Confirmed diagnosis of SHOX Syndrome
Epiphyses: Confirmation of open growth plates if age 10 and older

For short stature in children with Noonan Syndrome:
 ❖ **Norditropin®**

Height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Growth Velocity: < 5 cm/year
Bone age: Minimum of one year behind chronological age
Diagnosis: Confirmed diagnosis of Noonan Syndrome
Epiphyses: Confirmation of open growth plates if age 10 and older

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

For growth failure associated with chronic renal failure up to the time of transplantation:

❖ **Nutropin AQ®**

Renal function: Documentation of chronic renal insufficiency (glomerular filtration rate < 30 mL/min/1.73m²), up to the time of renal transplant
Height: ≥ 2 standard deviations (SD) below the mean for age and gender or less than the 5th percentile for age and gender
Growth Velocity: < 5 cm/year
Bone age: Minimum of one year behind chronological age
Epiphyses: Confirmation of open growth plates if age 10 and older
 Confirmation that existing metabolic derangements such as malnutrition, zinc deficiency, and secondary hyperparathyroidism have been corrected prior to initiation of GH treatment

❖ **Discontinuation of growth hormone therapy in children:**

- Expected final adult height has been reached; or
- If there is a poor response to treatment, generally defined as an increase in growth velocity of less than 50 % from baseline, in the 1st year of therapy; or
- Increase in height velocity is less than 2 cm total growth in 1 year of therapy; or
- **If age 10 or older, confirmation that the growth (epiphyseal) plates are closed (evidenced by x-ray) – linear growth can no longer occur in patients with epiphyseal closure; or**
- There are persistent and uncorrectable problems with adherence to treatment

❖ **Criteria for continuation of growth hormone therapy in children:**

- FDA approved diagnosis
- Prescribed by an endocrinologist, pediatric endocrinologist or pediatric nephrologist
- Growth velocity ≥ 2.5cm/year **AND**
- If age 10 and older confirmation that the bone age is less than 16 years in males; 14 years in females (indicated in x-ray of fingers, hands, or wrists) **AND**
- If age 10 or older, confirmation that the growth (epiphyseal) plates must be open (evidenced by x-ray) – linear growth can no longer occur in patients with epiphyseal closure

DOSING AND ADMINISTRATION:

- Refer to product labeling at <https://www.accessdata.fda.gov/scripts/cder/daf/>

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

GROWTH HORMONE TREATMENT IN ADULTS

Product Name	FDA Approved Indication	Comments
PREFERRED		
Short-Acting Agents		
Genotropin®, Norditropin®	Growth Hormone deficiency	N/A
Long-Acting Agents		
N/A		
NON-PREFERRED		
Short-Acting Agents		
Humatrop® [®] , Zomacton® [®] Nutropin/Nutropin AQ® [®] , Omnitrope® [®] , Saizen® [®]	Growth Hormone Deficiency	Refer to preferred short-acting agents
Long-Acting Agents		
Sogroya® [®]	Growth Hormone Deficiency	Must have documented trial and failure on 2 short-acting agents.

PRADER WILLI

- Growth hormone therapy is not approved in Prader Willi unless the beneficiary meets the growth hormone deficiency criteria for adults.

REVIEW CRITERIA FOR ADULTS:

- Must have approved diagnosis (see chart above for requested medication).
- The prescriber of the requested growth hormone must be an endocrinologist or gastroenterologist (for a diagnosis of short bowel syndrome).
- Patients with childhood-onset growth hormone deficiency (COGHD) previously treated with GH replacement in childhood should be retested after final height is achieved and GH therapy discontinued for at least 3 months to ascertain their GH status before considering restarting GH therapy (**at the reduced dose level recommended for growth hormone deficient adults**). (A repeat stimulation test may be required at the beginning of the next age increment in which a variation of IGF-1 occurs).
- For childhood GH treatment of conditions other than GHD, such as Turner’s syndrome and idiopathic short stature, there is no proven benefit to continuing GH treatment in adulthood.
- A negative response to a standard growth hormone stimulation test is a maximum peak of < 5 ng/ml, when measured by radioimmunoassay (RIA) (polyclonal antibody) or < 2.5 ng/ml when measured by immunoradiometric assay (monoclonal antibody).

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

- The preferred stimulation test agent is the Insulin Tolerance Test (ITT). Alternative provocative tests may be used in patients with contraindication to ITT. Other alternatives include glucagon, and rarely the arginine test alone. The glucagon stimulation test is associated with good performance and great diagnostic accuracy for GHD diagnosis:
 - *If a single agent test (arginine) is used there may be a requirement for a second stimulation test depending on the IGF-1. If the IGF-1 is subnormal with the presentation of a hypothalamic disorder(s) then one stimulation test would be required. However, if the IGF-1 is normal with hypothalamic pituitary disorder(s) then two stimulation tests may be required.*
 - *ITT is contraindicated in cases with coronary artery disease or seizures, abnormal EKG with history of ischemic heart disease or cardiovascular disease, and not advised for those > age 60.*
- Levodopa and Clonidine are not adequate agents for adult testing.
- The practitioner must correct for TSH deficiency prior to completing a stimulation test.
- A Growth Hormone stimulation test is not required when there is documented deficiencies of 3-4 pituitary hormones or documented deficiency of two pituitary hormones and IGF-1 < 84ng/ml. The anterior pituitary hormone deficiencies accepted for this exception to stimulation testing include: FSH and/or LH (subnormal results in both FSH and LH, simultaneously, would count as one deficiency), TSH, ACTH, and arginine vasopressin (AVP).
- Low IGF-1 alone is not an indicator of growth hormone deficiency.
- For diagnosis of short bowel syndrome, the prescriber must submit documentation to verify the diagnosis and the use of specialized nutrition support such as a high carbohydrate, low fat diet, enteral feedings, parenteral nutrition, fluid, and micronutrient supplements. Zorbtive® therapy is indicated under these conditions.
 - NOTE: Changes to concomitant medications should be avoided during Zorbtive® therapy.
 - **Subcutaneous dosage (Zorbtive® only):**
Adults and the elderly: 0.1 mg/kg SC once daily for 4 weeks. Do not exceed a maximum of 8 mg/day. Dosage selection for the elderly should usually start at the lower end of the dosage range. In clinical trials, Zorbtive® (plus a specialized oral diet without glutamine) vs. diet alone significantly decreased the total amount of intravenous parenteral nutrition (TPN) by 2.1L/week. The addition of glutamine to the diet/Zorbtive® group resulted in a significant decrease in IPN of 3.9 L/week. Other clinical reports have also documented a reduction in TPN usage.

DOSING AND ADMINISTRATION:

- Refer to product labeling at <https://www.accessdata.fda.gov/scripts/cder/daf/>

References:

1. American Academy of Pediatrics. Considerations related to the use of recombinant human growth hormone in children. American Academy of Pediatrics Committee on Drugs and Committee on Bioethics. Peds. 1997;99(1):122-129. Retrieved November 16, 2012, from: <http://pediatrics.aappublications.org/content/99/1/122.full>

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

2. American Association of Clinical Endocrinologists Medical Guidelines for Clinical Practice for Growth Hormone Use in Growth Hormone-Deficient Adults and Transition Patients-2009 Update. Retrieved November 9, 2012, from: <https://www.aace.com/files/growth-hormone-guidelines.pdf>
3. Barker AN, ed. LabPLUS Electronic Handbook. Auckland District Health Board. Auckland, NZ: LabPLUS; July 16, 2003. Available at: <http://www.adhb.govt.nz/LabPlusHandbook/>. Accessed August 26, 2003.
4. Blum WF, Crowe BJ, Quigley CA, Jung H, Cao D, Ross JL, Braun L, Rappold G. Growth hormone is effective in treatment of short stature associated with short stature homeobox-containing gene deficiency: two-year results of a randomized, controlled, multicenter trial. *J Clin Endocrinol Metab.* 2007 Jan;92(1):219-28. [PubMed]. Retrieved November 9, 2012, from: <http://jcem.endojournals.org/content/92/1/219.long>
5. Clayton PE, Cianfarani S, Czernichow P, et. al. Management of the child born small for gestational age through to adulthood: a consensus statement of the International Societies of Pediatric Endocrinology and the Growth Hormone Research Society. *J Clin Endocrinol Metab.* 2007; 92:804-810. Retrieved February 21, 2013, from: <http://www.ghresearchsociety.org/GRS%20consensus.htm>
6. Clemmons D. Consensus statement on the standardization and evaluation of growth hormone and insulin-like growth factor assays: a summary of the Growth Hormone Research Society. *Clinical Chem.* 2011;57(4):555-559. Retrieved February 21, 2013, from: <http://www.ghresearchsociety.org/GRS%20consensus.htm>
7. Cohen P, Rogol AD, Deal CL, Saenger P, et al. Consensus statement on the diagnosis and treatment of children with idiopathic short stature: a summary of the Growth Hormone Research Society, the Lawson Wilkins Pediatric Endocrine Society, and the European Society for Paediatric Endocrinology Workshop. *J Clin Endocrinol Metab.* 2008;93:4210–4217. [PubMed]. Retrieved November 9, 2012, from: <http://jcem.endojournals.org/content/93/11/4210.long>
8. de Zegher F. Small as Fetus and Short as Child: From Endogenous to Exogenous Growth Hormone. *Journal of Clinical Endocrinology & Metabolism.* 1997; 82(7):2021-2026.
9. Shalitin S. Children Born Small for Gestational Age: Growth Patterns, Growth Hormone Treatment and Long-Term Sequelae. *IMAJ.* 2003; 5:877-882.
10. Genotropin [package insert]. Kalamazoo, MI; Pharmacia and Upjohn Co/Pfizer; March 2011. Retrieved November 16, 2012, from: <http://dailymed.nlm.nih.gov/dailymed/drugInfo.cfm?id=45466>
11. Grimberg A, et al. Guidelines for Growth Hormone and Insulin-Like Growth Factor-I Treatment in Children and Adolescents: Growth Hormone Deficiency, Idiopathic Short Stature, and Primary Insulin-Like Growth Factor-I Deficiency. *Horm Res Paediatr.* 2016; 86:361-397.
12. Growth Hormone Research S. Consensus guidelines for the diagnosis and treatment of growth hormone (GH) deficiency in childhood and adolescence: summary statement of the GH Research Society. GH Research Society. *The Journal of clinical endocrinology and metabolism* 2000; 85:3990-3.
13. Growth Hormone Task Force of the American Association of Clinical Endocrinologists and the American College of Endocrinology. AACE medical guidelines for clinical practice for growth hormone use in adults and children, Jan/Feb 2003. Retrieved November 9, 2012, from: <http://misc.medscape.com/pi/editorial/pguidelines/aace/hgh.pdf>
14. Pedicelli S, Peschiaroli E, Violi E, Cianfarani S. Controversies in the definition and treatment of idiopathic short stature (ISS). *J Clin Res Pediatr Endocrinol.* 2009 March; 1(3): 105–115. [PubMed]. Retrieved November 9, 2012, from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3005647/>
15. Richmond EJ, Rogol AD. Growth hormone deficiency in children. *Pituitary* 2008; 11:115-20.
16. Sizonenko PC, Clayton PE, Cohen P, Hintz RL, Tanaka T, Laron Z. Diagnosis and management of growth hormone deficiency in childhood and adolescence. Part 1: diagnosis of growth hormone deficiency. *Growth hormone & IGF*

Division: Pharmacy Policy	Subject: Prior Authorization Criteria
Original Development Date: Original Effective Date: Revision Date:	June 8, 2012; April 4, 2013; July 8, 2013, February 25, 2015, June 11, 2015, January 21, 2016, February 12, 2016, May 19, 2016, July 5, 2016, March 7, 2017, August 18, 2017, March 6, 2018, July 9, 2020, September 23, 2020, October 15, 2020, September 22, 2021, January 7, 2022, July 26, 2022, September 8, 2022, January 26, 2023, February 9, 2024, February 20, 2024

research: official journal of the Growth Hormone Research Society and the International IGF Research Society. 2001;11(3):137-165.

17. Southern Reh C, Geffner M. Somatropin in the treatment of growth hormone deficiency and Turner syndrome in pediatric patients: a review. Clin Pharmacol. 2010; 2: 111–122. [PubMed]. Retrieved November 9, 2012, from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3262362/>
18. Tanner JM, Davies PSW. Clinical longitudinal standards for height and height velocity for North American children. J Pediatr. 1985;107:317-329.
19. Van Pareren Y, et al. Final Height in Girls with Turner Syndrome after Long-Term Growth Hormone Treatment in Three Dosages and Low Dose Estrogens. J Clin Endocrinol Metab. 2003; 88(3):1119–1125.
20. Wilson TA, Rose SR, Cohen P, Rogol AD, Backeljauw P, Brown R, Hardin DS, Kemp SF, Lawson M, Radovick S, Rosenthal SM, Silverman L, Speiser P. Update of guidelines for the use of growth hormone in children: The lawson wilkins pediatric endocrinology society drug and therapeutics committee. Journal of Pediatrics. 2003;143(4):415-421.